


The World of Webb, seeing through the Eyes of Einstein

Rogier Windhorst (ASU) — JWST Interdisciplinary Scientist

+JWST PEARLS team: T. Carleton, S. Cohen, R. Jansen, P. Kamieneski, T. Acharya, H. Archer, J. Berkheimer, D. Carter, N. Foo, R. Honor, D. Kramer, T. McCabe, I. McIntyre, R. O'Brien, R. Ortiz, J. Summers, S. Tompkins, C. Conselice, J. Diego, S. Driver, J. D'Silva, B. Frye, H. Yan, D. Coe, N. Grogin, W. Keel, A. Koekemoer, M. Marshall, N. Pirzkal, A. Robotham, R. Ryan Jr., C. Willmer + 100 more scientists over 18 time-zones



ASU SESE Undergraduate Student Seminar, Camp SESE, Tontozona, AZ

Saturday Sept. 9, 2023. All presented materials are ITAR-cleared.



North Ecliptic Pole (NEP) Time Domain Field (TDF) from PEARLS project:

(PEARLS = Prime Extragalactic Areas for Reionization and Lensing Science; Windhorst et al. 2023, *Astron. J.*, 165, 13; astro-ph/2209.04119)

- The NEP TDF is unique: Webb can observe it 365 days per year!
- (Old SED) tidal tails everywhere. Abundance of red (dusty) spirals
- $\lesssim 1\%$ of objects variable: Supernovae and accretion black-holes

Outline

- (1) Update on the James Webb Space Telescope (JWST), 2023.
- (2) Webb's first images: the "Cosmic Circle of Life"
- (3) Summary and Conclusions
- (4) What Hubble has done: Galaxy Assembly & SMBH Growth
 - (5) How can JWST measure Star-formation & Earth-like exoplanets?



WARNING:

Asking NASA for Hubble images is like drinking from a fire-hydrant,
and asking NASA for Webb images is like taking a sip from Niagara Falls!

Children: Please don't do this at home!! :)



Edwin P. Hubble (1889–1953) — Carnegie astronomer




James E. Webb (1906–1992) — Second NASA Administrator


Hubble: Concept in 1970's; Made in 1980's; Operational 1990– \gtrsim 2023?.

JWST: The infrared sequel to Hubble from 2021–2026 (–2031?).

**JWST primary
mirror**




**Hubble primary
mirror**



JWST $\simeq 2.5\times$ larger than Hubble, so at $\sim 2.5\times$ larger wavelengths:
JWST has the same resolution in the near-IR as Hubble in the optical.

(1) Update of the James Webb Space Telescope as of 2023



- A fully deployable 6.5 meter (25 m^2) segmented IR telescope for imaging and spectroscopy at $0.6\text{--}28 \mu\text{m}$ wavelength, launched Dec. 25, 2021.
- Nested array of sun-shields to keep its ambient temperature at 40 K, allowing faint imaging ($31.5 \text{ mag} \sim 1 \text{ FF}$ from Moon), and spectroscopy.

THE JAMES WEBB SPACE TELESCOPE

JWST LAUNCH


- LAUNCH VEHICLE IS AN ARIANE 5 ROCKET, SUPPLIED BY ESA
- SITE WILL BE THE ARIANESPACE'S ELA-3 LAUNCH COMPLEX NEAR KOUROU, FRENCH GUIANA



ARIANESPACE - ESA - NASA


- The JWST launch weight is $\lesssim 6500$ kg, and it was launched to L2 with an ESA Ariane-V launch vehicle from Kourou in French Guiana.

(1a) How did JWST travel to its L2 orbit?




- After launch on Dec. 25, 2021 with an ESA Ariane-V, JWST will orbit around the Earth–Sun Lagrange point L2, 1.5 million km from Earth.
- JWST can cover the whole sky in segments that move along with the Earth, observe $\gtrsim 70\%$ of the time, and send data back to Earth every day.

- (1b) How was JWST automatically deployed?




- During its two month journey to L2, JWST will be automatically deployed, its instruments will be cooled, and be inserted into an L2 orbit.
- The entire JWST deployment sequence is being tested several times on the ground — but only in 1-G: component and system tests in 2014–2019 at GSFC (MD), Northrop (CA), and JSC (Houston).
- Component fabrication, testing, & system integration: 18 out of 18 flight mirrors done in 2015, and meet the 40K specifications (2017).

Actuators for 6 degrees of freedom rigid body motion



Lightweighted
Beryllium Mirror

Actuator for radius
of curvature adjustment



Actuator
development
unit


Active mirror segment support through "hexapods", similar to Keck.
Redundant & doubly-redundant mechanisms, quite forgiving against failures.




JWST Hardware Status



Primary Mirror Segment



Aft Optics System



PM Flight Backplane



Tertiary Mirror



Fine Steering Mirror

Secondary Mirror Pathfinder Strut



ISIM Flight Bench



Secondary Mirror Hexapod



Secondary Mirror



Membrane Mgmt




Pathfinder Membrane




Spacecraft computer Test Unit







Mid-boom Test



2021: 100% of launch mass designed and built (100% weighed).



JWST Hardware Progress




JWST remains on track for an October 2018 launch within its replan budget guidelines

29

July 2014: ● Secondary Mirror Support deployment successfully tested.
2015: ● Engineering sunshield successfully deployed at Northrop (CA).


Much progress has been made in OTE integration



Where we were at last month's call

Current: all 18 PMSAs installed, liquid-shim-cured, & metrologized. Alignments meet specifications, and actuator motions verified

Big milestone!



8 February 2016 JWST Monthly Telecon 8

JWST lifetime: Requirement: 5 yrs; Goal: 10 yrs; Propellant: 20⁺ yrs!




NASA team-work to take JWST mirror covers off!



JWST being tilted into the right position





Webb mirrors finally mounted and ready!




JWST stowed for further instrument mounting




All Instruments Integrated



JWST: A Product of the Nation






European Consortium Who & Where




- JWST hardware made in 27 US States: 100% of launch-mass finished.
- Ariane V Launch & NIRSpec provided by ESA; & MIRI by ESA & JPL.
- JWST Fine Guider Sensor + NIRISS provided by Canadian Space Agency.
- JWST NIRCams made by UofA and Lockheed.


Astronomy Scene



Metal Mask/Fixed Slit



Shutter Mask






April 2017: Last portrait of JWST at Goddard Space Flight Center (MD).



SMSS Deployment Sequence (1)




190812 JWST Monthly Telecon 8

July 2019: Full 1-G deployment of JWST secondary mirror (SM) .



SMSS Deployment Sequence (2)




190812 JWST Monthly Telecon 9

July 2019: Full 1-G deployment of JWST secondary mirror (SM) ..



SMSS Deployment Sequence (3)







190812 JWST Monthly Telecon 10


July 2019: Full 1-G deployment of JWST secondary mirror (SM) ...



May 2017: JWST in enclosure at Johnson Space Center in Houston.








Aug. 2019: JWST OTE+ISIM lowered into Sunshield+Spacecraft



Meet the JWST Observatory 1



See NASA Press Release here:

<https://www.nasa.gov/feature/goddard/2019/nasa-s-james-webb-space-telescope-has-been-assembled-for-the-first-time>

© 2019 JWST Monthly Telecon 11

August 2019: JWST OTE+ISIM integrated with Sunshield and Spacecraft!




Transport to the Large Acoustic Test Facility

Primary Mirror Wing

Unitized Pallet Structure

Contamination Tent

Secondary Mirror



En route through the Space Park, Credit: NGSS

Arriving at the LATF Airlock, Credit: NGSS

2009-14 JWST Monthly Telecon 12

Aug 2020: Transport of JWST into Northrop acoustic chamber



(beautiful)
**The James Webb
Space Telescope**
Stowed for Launch



210913 JWST Monthly Telecon 18

Sept. 2021: JWST ready and stowed for shipping to Kourou



Dec. 9, 2021: JWST transport in Kourou to Ariane Rocket Assembly Building



Webb is finally launched from Kourou on December 25, 2021!



Feb. 2022: Webb seen shortly after launch over Africa using the Ariane V camera.

PRIMARY MIRROR SELFIE






IMAGE ARRAY



Feb. 2022: Webb's first selfie (left) and First Light raw image (right).

COMPLETED SEGMENT ALIGNMENT

COMPLETED IMAGE STACKING



Webb's first segment alignment (left) and first image stack (right).


TELESCOPE ALIGNMENT EVALUATION IMAGE




March 16, 2022: Webb's first fully focused image publicly released !!
Note the plethora of faint galaxies — Webb's looking back in time!

<https://www.nasa.gov/press-release/nasa-s-webb-reaches-alignment-milestone-optics-working-successfully>

WEBB TELESCOPE IMAGE SHARPNESS CHECK



April 28, 2022: Webb's first fully focused images in all four instruments: a dense star field in the Large Magellanic Cloud in the South Ecliptic Pole! (NIRSpec: $1.1 \mu\text{m}$; NIRISS: $1.5 \mu\text{m}$; NIRCam: $2.0 \mu\text{m}$; MIRI $7.7 \mu\text{m}$).




SPITZER IRAC 8.0 μ



WEBB MIRI 7.7 μ


May 9, 2022: Webb's 7.7. μ m MIRI image compared to Spitzer 8.0 μ m:
Same dense star field in the Large Magellanic Cloud in the South Ecliptic Pole

<https://blogs.nasa.gov/webb/2022/05/09/miris-sharper-view-hints-at-new-possibilities-for-science/>



Webb first NIRSpec near-IR spectra of ~ 100 faint stars near Galactic Center

Webb can take spectra of many 1000's of faint galaxies revealing their distances and chemical composition.



Hubble WFPC2 Eagle Nebula (1995) compared to JWST NIRCам (2022):


- The cradle of cosmic star-formation: NIRCам peers through the dust!
- The 1995 Hubble WFPC2 image (left) was made by Prof. Jeff Hester and Paul Scowen at ASU. It made it onto a US postage stamp!



Webb's MIRI shows the hauntingly beautiful cosmic dust pillars (8–15 μm)

JAMES WEBB SPACE TELESCOPE

TARANTULA NEBULA | NGC 2070






NIRCam Filters | F090W F200W F335M F444W

Tarantula Nebula “30 Doradus” in Large Magellanic Cloud (163,000 lyrs away)
Cradle of cosmic star-formation: massive stars trigger formation of sun-like stars



“Cosmic Cliffs” of star-formation in the Carina Nebula (NIR; 7600 light-years).

You will be witnessing the “Cosmic Circle of Life” ...



Where the action is: Composite of the Carina “Cosmic Cliffs”


- Red indicates molecular hydrogen outflowing from young forming stars.
- Young stars squirt out molecular jets that cause far-away bow-shocks.




Cosmic Cliffs of Star-formation in Carina Nebula (NIR+MIR).

Compared to optical+near-IR, mid-IR sees “Cradle of Cosmic Star-formation”


Deep inside the gas and dust, mid-IR reveals birth of young Sun-like stars.



NIRCam Near-Infrared



MIRI Mid-Infrared



MIRI and NIRCam


- JWST NIRCam+MIRI: Cosmic Cliff-like image of Orion's Trapezium:
- New stars are forming containing the carbon chain "Methyl Cation"



- Our birth, *e.g.*, : Protoplanetary “Hourglass Nebula” L1527 at 460 lyrs.
- A forming protostar with $\sim 30\%$ of Sun’s mass only 100,000 year old!
 - The protostar has surrounding accreting gas, and a circumstellar disk.
 - Eventually, L1527 will start shining as a star, and have its own planets.




- NIRCam+MIRI: ρ Ophiuchi dark cloud (stellar nursery closest to Earth):
- Cradle of star-formation contains Polycyclic Aromatic Hydrocarbons!



- NIRCam: newly forming stars Herbig-Haro 46/47 with jet-expelled material


FOMALHAUT



MIRI Filters | F2550W

JWST MIRI Coronagraph: Debris disk around nearby star Fomalhaut

- This is how the giant planets and terrestrial planets formed around our Sun




Dusty debris disk around red dwarf star AU Mic at 32 light-years.

- NIRCam's Coronagraph blocks the central star-light.
- Debris disk visible for 5–60 AU, *i.e.*, slightly larger than Solar System.


Mars

James Webb Space Telescope
NIRCam - September 5, 2022




Simulated Mars image with base maps
from NASA and MOLA data

NASA, ESA, CSA, STScI, MARS JWST/GTO team




NIRCam brightness at 2.1 microns



NIRCam brightness at 4.3 microns

Mars' surface with NIRCam: From "hot" to "cold" in the infrared!




Mars atmosphere NIRSpec spectrum: Plenty of Carbon Dioxide ...
but the search is much harder for Water vapor and Carbon Monoxide



Aug. 2022: JWST NIRcam image of the planet Jupiter: it has beautiful aurorae at its North and South pole — very strong magnetic field!
The Great “Red” Spot: A giant 4-century storm $2\times$ Earth’s diameter.

Saturn
JWST NIRCам F323N
June 25, 2023




JWST NIRCам: Our own planet Saturn with its moons and rings:


- Planetary rings are “failed moon” due to its strong tidal forces.

Titan

November 4, 2022




lower atmosphere and clouds



atmosphere and surface


Saturn's moon Titan: JWST NIRCams medium-band and color-composite.

- Bright clouds are visible in the near-IR, and they move with time!
- Kraken is a methane sea — it rains methane on Titan!
- Belet is a darkly colored plane with sand dunes.



NIRCam: Our own planet Uranus with new Zeta ring (*i.e.*, a failed moon)

- Polar cap: warmest point on Uranus for half its 88-year orbit!



NIRCam family portrait of Neptune with 7 of its Moons

Moon Triton is brighter, since methane darkens Neptune's atmosphere




Closeup of planet Neptune with Webb's NIRCам


- Giant planets with (dim) rings more common those than without rings!

Star
HIP 65426

Exoplanet
HIP 65426 b




Webb 3–15 micron exoplanet images (10 Jupiter masses; 15 Myr young!)




- Hot exoplanet WASP-96b orbiting a Sun-like star (1150 light-years):
- Near-IR spectrum shows characteristic features of water (steam !).
- It has a temperature of 1000 F and is half Jupiter in mass.
- Webb will scan Earth-like exoplanets for building blocks of life.

HOT GAS GIANT EXOPLANET WASP-39 b ATMOSPHERE COMPOSITION


NIRISS | Single Object Slitless Spectroscopy




NIRCam F322W2



NIRSpec G395H




NIRSpec PRISM



WEBB
SPACE TELESCOPE

Hot exoplanet WASP-39b orbiting a Sun-like star (700 light-years):

- Near-IR spectrum shows characteristic features of water (steam !).
- It has a temperature of 1650 F and is about Saturn's mass.
- Complex — and poisonous — Sulfur and Silicates atmosphere.



Southern Ring Nebula (Near-IR+Mid-IR; 2500 light-years):


- You **are** witnessing the “Cosmic Circle of Life” here ...
- This is a Sun-like star expelling its outer layers in retirement ...
- It has exhausted its hydrogen and helium as nuclear fuel ...
and expanded to $100\times$ its current size, engulfing the Earth.



This is how our Sun *will* come to an end in 5 Billion years (near-IR).
“... for dust thou art, and unto dust shalt thou return” (Genesis 3:19).




From gas expelled by previous sun-like stars, new stars are born (mid-IR).
And thanks to the dust they expelled, new stars will form with planets ...



30 solar mass Wolf Rayet star WR124 shortly before it turns Supernova ...

- [Left] NIRCam and [Right] MIRI — both showing recent mass loss.
- Prelude stage to Supernova also releases a lot of (dusty) mass!

JAMES WEBB SPACE TELESCOPE
SUPERNOVA 1987A



NIRCam Filters | F150W F164N F200W F322N F405N F444W

- NIRCam: Remnants of Supernova 1987A seen in Large Magellanic Cloud
- Shells outflowing over the decades caused hour-glass shaped bubbles



JWST MIRI: Supernova Remnant Cassiopeia-A expelling dust




Stephan's Quintet: 4 colliding galaxies (40 M-lyr; left spiral is foreground).

- These major “Cosmic Trainwrecks” are much more common in the past.
- Sun-like stars formed in aftermath of minor “Cosmic Fender-benders”.



- Stephan's Quintet: 4 colliding galaxies at 40 million light-years (Mid-IR):
- Mid-IR shows molecular gas being pulled out during collision.
 - Gravity from collision in top galaxy feeds the Beast: central black hole!



Hubble



Webb

Galactic Merger: Hubble vs. Webb — a veritable cosmic train wreck!

- [Left]: Hubble sees the young star-forming regions and dust.
- [Right]: Webb sees also the warm dust in the infrared (orange filaments).

(First revealed by Vice President Harris & French President Macron; Dec. 2022)




NGC1433 a galaxy with dusty spiral arms at 48 million light-years



NGC7496 a galaxy with dusty spiral arms at 24 million light-years:

- Inner spiral arms feed the central monster (black hole!)



Don't feed the animals: NGC7469 a spiral galaxy at 220 million light-years:

- It has a supermassive black hole feasting on the in-falling gas!
- In area surrounding the SMBH, gas is expelled at very high speeds, and stars are forming in ambient cooler gas.




JWST NIRcam+MIRI: nearby actively star-forming galaxy Arp 220:


- Copious amounts of inflowing gas and dust feed the central monster!




Spiral Galaxy overlapping Elliptical: Tracing cosmic dust (Keel⁺ 22) ...




... and the elliptical also lenses a galaxy seen 2 Byrs after Big Bang!



LEDA-2046648: a beautiful galaxy pair observed with NIRISS 1 Blyr away



LEDA-2046648: Andromeda will collide will Milky Way like this in 4-5 Byrs.



Will this ever happen to our own Galaxy?

YES! Hubble showed no lateral motion of Andromeda:


Approaches at -110 km/s.

Hence, Andromeda will merge with Milky Way!


The two blackholes (10^6 – 10^7 suns) will also merge!

Not to worry: only 4–5 Byr from today!


Illustration Sequence of the Milky Way
and Andromeda Galaxy Colliding



July 11, 2022: 12-hr Webb Deep Field on galaxy cluster SMACS 0723





Hubble image of SMACS 0723 – Webb sees the dawn of galaxy formation!



Cluster RXJ2129 with triply lensed Supernova at 2.9 billion lyrs distance

- SN only seen in middle panel sampling the earliest observation

<https://esawebb.org/images/potm2302a/>




Massive lensing cluster Abell 2744:

Over 10^{15} solar masses seen 4 billion years ago:


Its gravity lenses 5 young galaxies at redshift $z \simeq 7.88$,

i.e., magnifying objects seen 13 billion years ago.

Webb is looking back to 650 million years after Big Bang!



Cluster MACS0647 triply lensed a galaxy 0.4 Byrs after BB! (Hsiao, Coe⁺ 22)




NIRCam: Lensing cluster WHL0137-08 with highly lensed arc at $z=6.2$

- Earendel: a highly magnified (double-)star seen in the first billion years after the Big Bang!



JWST image of most luminous far-IR Planck cluster G165 at $z=0.35$ found:
Highest- z SN-Ia: 3 epochs, 9 points at $z=1.78 \rightarrow H_0$? (Frye, Foo⁺ 2023).




Monster cluster El Gordo distorts distant galaxies into “pencils” (Diego⁺22)





and El Gordo makes a super-lens “El Anzuelo” — Einstein’s fishhook!

17 hr JWST on Hubble Frontier Field cluster MACS0416 ($z=0.397$; 4 Blyr)




JWST: Lensed Dusty sources behind El Gordo in first few Byrs (P. Kamieneski⁺; astro-ph/2303.05054):




13 filter Hubble UltraDeep Field:

UV-Blue emphasized.




○ $z=7-8$, ○ $z=9$, ○ $z=10-12$.


Hubble UltraDeep Field: Red-IR emphasized.



13 filter Hubble UltraDeep Field:

False-color image: sky is half covered by galaxies.







Conclusion: JWST First Light strategy must consider three aspects:

- (1) The earliest objects (first 0.5 Byr) are very rare and hard to find.
- (2) Cannot-see-the-forest-for-the-trees effect [“Confusion” limit]:
Background objects blend into foreground objects because of their density.
- (3) House-of-mirrors effect [“Gravitational Amplification”]:
 - Lensing is needed to see what Einstein thought was impossible to observe!

(3) Summary and Conclusions

(1) Webb was successfully built, tested and finally launched in Dec. 2021.

(2) Webb was designed to map the epochs of First Light, Galaxy Assembly & Super Massive Black Hole-growth in detail:

- Formation of the first stars and star-clusters after 0.2 Byr.
- How galaxies formed and evolved over 13.5 Billion years.

(3) Webb's first images trace the "Cosmic Circle of Life":

- Formation and evolution of stars and dust over cosmic time.
- How dust helped form exoplanets and building blocks for life.

(4) Webb will have a major impact on astrophysics this decade:

- IR sequel to HST starting 2022: Training next generation researchers.

SPARE CHARTS

● References and other sources of material

Talk: http://www.asu.edu/clas/hst/www/jwst/asu_campSESE23_hstjwst.pdf Data: <https://sites.google.com/view/jwstpearls>

Carleton, T., Windhorst, R. A., O'Brien, R., et al. 2022, AJ, 164, 170 (astro-ph/2205.06347)

Carleton, T., Cohen, S. H., Frye, B., et al. 2023, ApJ, 953, 83 (astro-ph/2303.04726)

Cheng, C., Huang, J.-S., Smail, I., et al. 2023, ApJ, 942, L19 (astro-ph/2210.08163)

Diego, J. M., Meena, A. K., Adams, N. J., et al. 2023, A&A, 672, A3 (astro-ph/2210.06514)

Duncan, K. J., Windhorst, R. A., et al. 2023, MNRAS, 522, 4548–4564 (astro-ph/2212.09769)

Ferreira, L., Adams, N., Conselice, C. J., et al. 2022, ApJL, 938, L2 (astro-ph/2207.09428)

Frye, B. L., Pascale, M., Foo, N., et al. 2023, ApJ, 952, 81 (astro-ph/2303.03556)

Kamieneski, P. S., Frye, B. L., Pascale, M., et al. 2023, ApJ, in press (astro-ph/2303.05054)

Keel, W. C., Windhorst, R. A., Jansen, R. A., et al. 2023, AJ, 165, 166 (astro-ph/2208.14475)

Kramer, D. M., Carleton, T., Cohen, S. H., et al. 2022, ApJL, 940, L15 (astro-ph/2208.07218v2)

O'Brien, R., Carleton, T., Windhorst, R. et al. 2023, AJ, 165, 237 (astro-ph/2210.08010)

Polletta, M. del Carmen, Nonino, M., Frye, B., et al. 2023, A&AL, 675, L4 (astro-ph/2306.12385)

Summers, J., Windhorst, R. A., Cohen, S. H., et al. 2023, ApJ, resubmitted (astro-ph/2306.13037)

Windhorst, R., Timmes, F. X., Wyithe, J. S. B., et al. 2018, ApJS, 234, 41 (astro-ph/1801.03584)

Windhorst, R. A., Carleton, T., O'Brien, R., et al. 2022, AJ, 164, 141 (astro-ph/2205.06214)

Windhorst, R. A., Cohen, S. H., Jansen, R. A., et al. 2023, AJ, 165, 13 (astro-ph/2209.04119)

Yan, H., Cohen, S. H., Windhorst, R. A., et al. 2023, ApJL, 942, L8 (astro-ph/2209.04092)

<https://hubblesite.org/contents/news-releases/2022/news-2022-050>

<https://blogs.nasa.gov/webb/2022/10/05/webb-hubble-team-up-to-trace-interstellar-dust-within-a-galactic-pair/>

<https://blogs.nasa.gov/webb/2022/12/14/webb-glimpses-field-of-extragalactic-pearls-studded-with-galactic-diamonds/>

<https://esawebb.org/images/pearls1/zoomable/>

<https://webbtelescope.org/contents/news-releases/2023/news-2023-119>

<https://news.asu.edu/20230801-jwsts-gravitational-lens-reveals-distant-objects-behind-el-gordo-galaxy-cluster>


What the Scientists See:



What the Project Manager Sees:




The Happy Balance



Any (space) mission is a balance between what science demands, what technology can do, and what budget & schedule allows ... (courtesy Prof. R. Ellis).

3) How can Webb measure First Light: What to expect in (Ultra)Deep Fields?




- Can't beat redshift: to see First Light, must observe near-mid IR.
- ⇒ This is why JWST needs NIRCam at 0.8–5 μm and MIRI at 5–28 μm .

SDSS J165202.64+172852.3

MOTIONS OF GAS AROUND AN EXTREMELY RED QUASAR

Hubble ACS + WFC3 Imaging

Webb NIRSpec IFU Spectroscopy



WEBB
SPACE TELESCOPE


- NIRSpec spectral cube of a luminous quasar seen 2.2 Byrs after Big Bang. Colors indicate 3 companion galaxies falling into the quasar host galaxy.
- In the first 2 billion years big galaxies were swallowing little ones!

(4) What Hubble has done: Panchromatic High-Throughput Camera



HST WFC3 and its **IR channel**: a critical pathfinder for JWST science.

(4) Hubble WFC3: Measuring Galaxy Assembly and SMBH Growth?



10 filters with Hubble WFC3 & ACS reaching $AB=26.5-27.0$ mag over 40 arcmin² with $0.07-0.15''$ images from $0.2-1.7\mu\text{m}$ (UVUBVizYJH).

JWST adds $0.05-0.2''$ FWHM imaging to $AB\approx 31.5$ mag (1 FF) at $1-5\mu\text{m}$, with $0.2-1.2''$ images at $5-29\mu\text{m}$, tracing young+old stars & dust.

Black Hole growth — Waves that happen in Nature: 1) Sounds Waves:





In solids: Earthquakes



In liquids: Surf!





In gasses: Sound



2) Electromagnetic Waves

3) In space-time: Gravity Waves

Sept. 2015: LIGO added Gravity Waves as a new way to observe Nature!





(1) LIGO first observed Gravitational Waves on Sept. 14, 2015.

(2) These were caused by two merging ($29+36 M_{\odot}$) black holes about 1 Gyr ago!

● $E=Mc^2$: $3 M_{\odot}$ was converted to energy in a fraction of a second!

Visible

Infrared




30 Doradus Nebula and Star Cluster
Hubble Space Telescope ■ WFC3/UVIS/IR


NASA, ESA, F. Paresce (INAF-IASF, Italy), and the WFC3 Science Oversight Committee

STScI-PRC09-32b

30 Doradus: Giant young star-cluster in Large Magellanic Cloud (150,000 ly), triggering birth of Sun-like stars (and surrounding debris disks).



Ordinary massive stars ($10\text{--}30 M_{\odot}$) leave modest black holes ($\sim 3\text{--}10 M_{\odot}$).



Ordinary massive stars ($10\text{--}30 M_{\odot}$) leave modest black holes ($\sim 3\text{--}10 M_{\odot}$).

Conclusion 1: Most low-mass black holes today are small, slow eaters:



- 29–36 M_{\odot} blackholes may be leftover from First Stars (first 500 Myr).
- Likely too massive to be leftover from ordinary Supernova explosions, ...
- How come only now seen merging by LIGO (12.5 Byr after BB)?
- They were likely not fast & efficient eaters, but slow and messy ...

Elliptical galaxy M87 with Active Galactic Nucleus (AGN) and relativistic jet:



The danger of having Quasar-like devices too close to home ...
They are **EXTREMELY** bright sources if viewed "down-the-pipe".
~0.5% of the baryonic mass, but produce most of the photons!






Figure 15. Averages of the three fiducial images of M87 for each of the four observed days after restoring each to an equivalent resolution, as in Figure 14. The indicated beam is $20 \mu\text{as}$ (i.e., that of DIFMAP, which is always the largest of the three individual beams).


2019 discovery of Black Hole Shadow in M87 by Event Horizon Telescope:
M87 at 55 Mlyr distance has a black hole mass of $\sim 6.5 \times 10^9 M_{\odot}$!


Centaurus A
NGC 5128
HST WFC3/UVIS

F225W+F336W+F438W
F487N H β
F502N [O III]
F547M y
F657N H α + [N II]
F673N [S II]
F814W I

3000 light-years
1400 parsecs


56''





Blue=X-rays; White=Optical; Orange=Radio

- Quasars: Centers of galaxies with feeding supermassive blackholes:




- Hubble IR-images of the most luminous Quasar known in the universe.
- Seen at redshift 6.42 (universe $7.42\times$ smaller than today), 900 Myr old!
- Contains 10^{14} solar luminosities within a region as small as Pluto's orbit!
- A feeding monster blackhole ($>3\times 10^9$ solar mass) 900 Myr after BB!

Conclusion 2: Supermassive black holes started early & were very rapid eaters:




- Massive galaxies today contain a super-massive blackhole, no exceptions!
- Masses $\sim 3 \times 10^9$ solar, leftover from the First Stars (first 500 Myr)?
- Must have fed enormously rapidly in the first 1 Byr after the Big Bang.
- Were eating *cat*-astrophically (and secretly) until they ran out of food ...
- JWST can image the First Quasars to $z \gtrsim 10$ (*if* we can find them).

(5) How can JWST measure Star-Formation and Earth-like exoplanets?



NGC 3603: Young star-cluster triggering star-birth in “Pillars of Creation”


Pillar and Stellar Jet in the Carina Nebula
HST WFC3/UVIS



HST WFC3/IR




Fomalhaut
HST ACS/HRC



HST/ACS Coronagraph imaging of planetary debris disk around Fomalhaut:
First direct imaging of a moving planet forming around a nearby star!

JWST can find such planets much closer in for much farther stars.

Exoplanet HR 8799 System



HST/NICMOS

HST/NICMOS with additional processing

HR 8799 planet orbits

Relative size of
Neptune's orbit

NASA, ESA, and R. Soummer (STScI)

STScI-PRC11-29


HST/NICMOS imaging of planetary system around the (carefully subtracted) star HR 8799: Direct imaging of planets around a nearby star.

Press release: <http://hubblesite.org/newscenter/archive/releases/2011/29/>

JWST can find such planets much closer in for much farther-away stars.

Schematic of Transit and Eclipse Science

Seager & Deming (2010, ARAA, 48, 631)



Transit

Measure size of planet 10^{-2}

See starlight transmitted through planet atmosphere 10^{-4}


Eclipse

Planet thermal emission appears and disappears 10^{-3}

Learn about atmospheric circulation from thermal phase curves


6

Timeline of a Transit Observation




13

JWST can do very precise photometry of transiting Earth-like exoplanets.



JWST IR spectra can find water and CO₂ in (super-)Earth-like exoplanets.

Transit Spectrum of Habitable "Ocean Planet"



JWST IR spectra can find water and CO₂ in transiting Earth-like exoplanets.