


# Beyond Hubble: From Exoplanets to the First Stars with the James Webb Space Telescope

---

**Rogier Windhorst (ASU) — JWST Interdisciplinary Scientist**

*Collaborators: S. Cohen, L. Jiang, R. Jansen (ASU), C. Conselice (UK), S. Driver (OZ), & H. Yan (U-MO)*

*(Ex) ASU Grads: N. Hathi, H. Kim, M. Mechtley, R. Ryan, M. Rutkowski, A. Straughn & K. Tamura*



*Colloquium at Rome University, Physics Department, Rome, Italy*

*Tuesday March 11, 2014. All presented materials are ITAR-cleared.*

# Outline

---

- (1) The Best of Hubble: Recent results from the Hubble Space Telescope (HST) and its Wide Field Camera 3 (WFC3).
- (2) Measuring Star-birth and Earth-like exoplanets
- (3) Measuring Galaxy Assembly and Supermassive Black-Hole Growth.
- (4) What is the James Webb Space Telescope (JWST)?
- (5) How can JWST measure the Epochs of First Light & Reionization?
- (6) Summary and Conclusions.



Edwin P. Hubble (1889–1953) — Carnegie astronomer




James E. Webb (1906–1992) — Second NASA Administrator


Hubble: Concept in 1970's; Made in 1980's; Operational 1990– $\gtrsim$ 2014.

JWST: The infrared sequel to Hubble from 2018–2023 (–2029?).

**JWST primary  
mirror**



**Hubble primary  
mirror**



JWST  $\simeq 2.5\times$  larger than Hubble, so at  $\sim 2.5\times$  larger wavelengths:  
JWST has the same resolution in the near-IR as Hubble in the optical.

# (1) The Best of Hubble: Recent results from the HST and its WFC3




# WFC3: Hubble's new Panchromatic High-Throughput Camera




HST WFC3 and its **IR channel**: a critical pathfinder for JWST science.

## Role of ACS in HST Post-SM4 Imaging Capability



ACS/WFC superior to WFC3 survey efficiency at visible-red wavelengths



WFC3/UVIS channel unprecedented UV–blue throughput & area:

- $QE \gtrsim 70\%$ ,  $4k \times 4k$  array of  $0''.04$  pixel,  $FOV \simeq 2'.67 \times 2'.67$ .

WFC3/IR channel unprecedented near–IR throughput & area:


- $QE \gtrsim 70\%$ ,  $1k \times 1k$  array of  $0''.13$  pixel,  $FOV \simeq 2'.25 \times 2'.25$ .

⇒ WFC3 opened major new parameter space for astrophysics in 2009:

WFC3 filters designed for star-formation and galaxy assembly at  $z \simeq 1-8$ :

- HST WFC3 and its IR channel a critical pathfinder for JWST science.

## (2) Measuring Star-birth and Earth-like exoplanets





NGC 3603: Young star-cluster triggering star-birth in “Pillars of Creation”



Visible

Infrared



**30 Doradus Nebula and Star Cluster**  
*Hubble Space Telescope* ■ WFC3/UVIS/IR

NASA, ESA, F. Paresce (INAF-IASF, Italy), and the WFC3 Science Oversight Committee


STScI-PRC09-32b

30 Doradus: Giant young star-cluster in Large Magellanic Cloud (150,000 ly), triggering birth of Sun-like stars (and surrounding debris disks).






Pillar and Stellar Jet in the Carina Nebula  
HST WFC3/UVIS



HST WFC3/IR



## Exoplanet HR 8799 System



NASA, ESA, and R. Soummer (STScI)

STScI-PRC11-29


HST/NICMOS imaging of planetary system around the (carefully subtracted) star HR 8799: Direct imaging of planets around a nearby star!

Press release: <http://hubblesite.org/newscenter/archive/releases/2011/29/>

**JWST can find such planets much closer in for much farther-away stars!**

## Schematic of Transit and Eclipse Science

Seager & Deming (2010, ARAA, 48, 631)




**Eclipse**  
Planet thermal emission appears and disappears  $10^{-3}$

**Transit**  
Measure size of planet  $10^{-2}$   
See starlight transmitted through planet atmosphere  $10^{-4}$

Learn about atmospheric circulation from thermal phase curves


6

## Timeline of a Transit Observation




13

JWST can do very precise photometry of transiting Earth-like exoplanets.



JWST IR spectra can find water and CO<sub>2</sub> in (super-)Earth-like exoplanets.


# Transit Spectrum of Habitable "Ocean Planet"



JWST IR spectra can find water and CO<sub>2</sub> in transiting Earth-like exoplanets.

### (3) HST turned the classical Hubble sequence upside down!


## Edwin Hubble's Classification Scheme



Who (when)	Cosmic Epoch	Ellipticals	Spirals	Irr's/mergers
Hubble (1920's)	$z=0$ (13.73 Gyr)	$\sim 40\%$	$\gtrsim 50\%$	$\lesssim 10\%$
HST (1990's)	$z \simeq 1-2$ (3-6 Gyr)	$\lesssim 15\%$	$\sim 30\%$	$\gtrsim 55\% !$




Spiral Galaxy M83  
HST WFC3/UVIS



F336W U  
F555W V  
F814W I  
F502N [O III]  
F657N H $\alpha$


5,000 light-years  
1,500 parsecs  
70''







HST Antenna galaxy: Prototype of high redshift, star-forming, major merger?




**Illustration Sequence of the Milky Way  
and Andromeda Galaxy Colliding**


NASA, ESA, Z. Levay and R. van der Marel (STScI), T. Hallas, and A. Mellinger ■ STScI-PRC12-20b

Merger of Andromeda galaxy (M31) with Milky Way about 4 Gyr from now.

# Interacting Galaxies

Hubble Space Telescope • ACS/WFC • WFPC2





“Tadpole” Galaxies in the Hubble Ultra Deep Field  
*Hubble Space Telescope* ■ ACS/WFC


NASA, ESA, A. Straughn, S. Cohen and R. Windhorst (Arizona State University), and the HUDF team (STScI)

STScI-PRC06-04

Merging galaxies constitute  $\lesssim 1\%$  of Hubble sequence TODAY (age  $\gtrsim 12.5$  Gyr).

Tadpole galaxies are early stage mergers, very common at  $z \gtrsim 2$  (age  $\lesssim 3$  Gyr).


**JWST will measure Galaxy Assembly to  $z \lesssim 20$  (cosmic age  $\gtrsim 0.2$  Gyr).**



HST/WFC3 & ACS reach  $AB=26.5-27.0$  mag ( $\sim 100$  fireflies from Moon) over  $0.1 \times$  full Moon area in 10 filters from  $0.2-2\mu\text{m}$  wavelength.

JWST has  $3 \times$  sharper imaging to  $AB \simeq 31.5$  mag ( $\sim 1$  firefly from Moon) at  $1-5\mu\text{m}$  wavelengths, tracing young and old stars + dust.

### (3) Measuring Galaxy Assembly & Supermassive Blackhole Growth





**Radio Galaxy 0313-192**  
**Hubble Space Telescope ACS WFC • Very Large Array**

NASA, NRAO/AUI/NSF and W. Keel (University of Alabama) • STScI-PRC03-04

Does galaxy assembly go hand-in-hand with supermassive blackhole growth?



Elliptical galaxy M87 with Active Galactic Nucleus (AGN) and relativistic jet:



The danger of having Quasar-like devices too close to home ...


Centaurus A  
NGC 5128  
HST WFC3/UVIS

F225W+F336W+F438W  
F487N H $\beta$   
F502N [O III]  
F547M  $\gamma$   
F657N H $\alpha$ + [N II]  
F673N [S II]  
F814W I

3000 light-years  
1400 parsecs  
56''




### (3) HST WFC3 observations of QSO host galaxies at $z \simeq 6$ (age $\lesssim 1$ Gyr)




- Careful contemporaneous orbital PSF-star subtraction: Removes most of “OTA spacecraft breathing” effects (Mechtley et al 2012, ApJL, 756, L38).
- PSF-star ( $AB \simeq 15$  mag) subtracts  $z=6.42$  QSO ( $AB \simeq 18.5$ ) nearly to the noise limit: NO host galaxy detected  $100\times$  fainter ( $AB \gtrsim 23.5$  at  $r \gtrsim 0.3$ ).

### (3) WFC3: First detection of one QSO Host Galaxy at $z \simeq 6$ (Giant merger?)



- Monte Carlo Markov-Chain of PSF-star + Sersic ML light-profile: (Mechtley, Jiang, Windhorst et al. 2013; Mechtley 2013, PhD):
- FIRST solid detection out of four  $z \simeq 6$  QSOs [3 more to be observed].
- One  $z \simeq 6$  QSO host galaxy: Giant merger morphology + tidal structure??
- Same J+H-band structure! Blue UV spectrum: Constrains dust.
- Starburst-like spectrum from rest-frame UV to far-IR ( $A_{FUV} \sim 1$  mag).
- $L(z \simeq 6 \text{ host galaxy}) \simeq 6 \times$  brighter than typical  $L^*$ : Monster!
- Quasar duty cycle could be  $\lesssim 10^{-2}$  ( $\lesssim 10$  Myrs): Eats like a beast!


## (4) What is the James Webb Space Telescope (JWST)?



Need young generation of students & scientists after 2018 ... It'll be worth it!

(RIGHT) Life-size JWST prototype on the Capitol Mall, May 2007 ...

## (4) What is the James Webb Space Telescope (JWST)?



- A fully deployable 6.5 meter ( $25 \text{ m}^2$ ) segmented IR telescope for imaging and spectroscopy at  $0.6\text{--}28 \mu\text{m}$  wavelength, to be launched in Fall 2018.
- Nested array of sun-shields to keep its ambient temperature at 40 K, allowing faint imaging (31.5 mag = firefly from Moon!) and spectroscopy.

# THE JAMES WEBB SPACE TELESCOPE

## JWST LAUNCH


- LAUNCH VEHICLE IS AN ARIANE 5 ROCKET, SUPPLIED BY ESA
- SITE WILL BE THE ARIANESPACE'S ELA-3 LAUNCH COMPLEX NEAR KOUROU, FRENCH GUIANA



ARIANESPACE - ESA - NASA

- The JWST launch weight will be  $\lesssim 6500$  kg, and it will be launched to L2 with an ESA Ariane-V launch vehicle from Kourou in French Guiana.


## (4a) How will JWST travel to its L2 orbit?



- After launch in 2018 with an ESA Ariane-V, JWST will orbit around the Earth–Sun Lagrange point L2, 1.5 million km from Earth.
- JWST can cover the whole sky in segments that move along with the Earth, observe  $\gtrsim 70\%$  of the time, and send data back to Earth every day.





- (4b) How will JWST be automatically deployed?



- During its two month journey to L2, JWST will be automatically deployed, its instruments will be cooled, and be inserted into an L2 orbit.
- The entire JWST deployment sequence will be tested several times on the ground — but only in 1-G: Component and system tests in Houston.
- Component fabrication, testing, & system integration is on schedule: 18 out of 18 flight mirrors completely done, and meet the 40K specifications!

# Actuators for 6 degrees of freedom rigid body motion



Actuator development unit

Lightweighted Beryllium Mirror

Actuator for radius of curvature adjustment


Active mirror segment support through "hexapods", similar to Keck.  
Redundant & doubly-redundant mechanisms, quite forgiving against failures.



# JWST Hardware Status



Primary Mirror Segment



Aft Optics System



PM Flight Backplane



Tertiary Mirror

Secondary Mirror Pathfinder Strut



Fine Steering Mirror



Secondary Mirror Hexapod



ISIM Flight Bench



Secondary Mirror



Membrane Mgmt




Pathfinder Membrane



Spacecraft computer Test Unit




Mid-boom Test



80% of launch mass designed and built as of Jan. 2014.


# Mirror Acceptance Testing







# Primary Mirror Composite



RMS: **23.2 nm**

PV: **515.5 nm**





# Mirror Status



- **15 flight primary mirrors and the flight secondary mirror are at GSFC in storage**
  - All spares were at GSFC in storage (SM spares, 3 PMSA spares)
    - 2 EDU mirrors sent back to Ball for gear motor rework
  - All flight gear motor refurbishment is complete
  - All flight mirrors will be at GSFC by end of year, needed in 2015




Jan 2014: All 18 flight mirrors now delivered to NASA GSFC (MD).




# Backplane Support Frame, Center Section, & Wings



- Center Section is complete
- Wings and cryo cycling is complete
- BSF assembly is complete
- Integration of the BSF to Center Section Complete
  - Cryo Cycling at MSFC XRCF complete



BSF and Center Section



BSF/Center Section coming out of

Jan 2014: Flight back-plane ready to receive mirrors in 2014.



# Telescope Assembly Ground Support Equipment



Ambient Optical Alignment Stand



Hardware has been installed at GSFC approximately 8 weeks ahead of schedule



March 2012 NAC Science Meeting



Landing a mirror onto backplane simulator




# Sunshield Template Membrane Work Completed

Templates Verify Design/Manufacturing Prior to Flight Build



- All Template Layers Completed
- Preparing for flight article manufacturing
- First two Flight Manufacturing Readiness Reviews Completed
- Membrane pull out test complete

Stringing Operations



Template Layers 3-5



Hole Tool Operations

Flight sunshield to be completed & tested by 2015 at Northrop (CA).




## Instrument Overview





### Fine Guidance Sensor (FGS)

- Ensures guide star availability with >95% probability at any point in the sky
- Includes Narrowband Imaging Tunable Filter
- Developed by Canadian Space Agency & COM DEV




### Near Infra-Red Camera (NIRCam)

- Detects first light galaxies and observes galaxy assembly sequence
- 0.6 to 5 microns
- Supports Wavefront Sensing & Control
- Developed by Univ. of AZ & LMATC



### Mid-Infra-Red Instrument (MIRI)

- Distinguishes first light objects; studies galaxy evolution; explores protostars & their environs
- Imaging and spectroscopy capability
- 5 to 27 microns
- Cooled to 7K by Cyro-cooler
- Combined European Consortium/JPL development




### Near Infra-Red Spectrograph (NIRSpec)


- Measures redshift, metallicity, star formation rate in first light galaxies
- 0.6 to 5 microns
- Simultaneous spectra of >100 objects
- Developed by ESA & EADS with NASA/ GSFC Detector & Microshutter Subsystems



# JWST: A Product of the Nation



## European Consortium Who & Where



- JWST hardware made in 27 US States:  $\approx 80\%$  of launch-mass finished.
- Ariane V Launch & NIRSpec provided by ESA; & MIRI by ESA & JPL.
- JWST Fine Guider Sensor + NIRISS provided by Canadian Space Agency.
- JWST NIRCам made by UofA and Lockheed.


Thank you, Italy, for your contributions to JWST!




Aug. 2013: Actual Flight ISIM (with MIRI and FGS) lowered into OSIM.





# TELESCOPE ARCHITECTURE





# OTE Testing – Chamber A at JSC




Notice people for scale

Will be the largest cryo vacuum test chamber in the world

OTIS: Largest TV chamber in world: will test whole JWST in 2015–2016.

# Implications of the WMAP year-9 & Planck13 results for JWST science:



HST/WFC3  $z \lesssim 7-9$  ← — → JWST  $z \simeq 8-25$

The year-9 WMAP data provided better foreground removal (Komatsu<sup>+</sup> 2011; Hinshaw<sup>+</sup> 2012; but see: Planck XVI 2013.)


⇒ First Light & Reionization occurred between these extremes:

- (1) Instantaneous at  $z \simeq 11.1 \pm 1.1$  ( $\tau = 0.089 \pm 0.013$ ), or:
- (2) Inhomogeneous & drawn out: starting at  $z \gtrsim 20$ , peaking at  $z \lesssim 11$ , ending at  $z \simeq 7$ . The implications for HST and JWST are:
- HST/ACS has covered  $z \lesssim 6$ , and WFC3 is covering  $z \lesssim 7-9$ .
- For First Light & Reionization, JWST will survey  $z \simeq 8$  to  $z \simeq 15-20$ .

Question: If Planck- $\tau \downarrow \lesssim 0.08$  (TBD, Planck14), then how many reionizers will JWST see at  $z \simeq 10-20$ ?




## (5a) How will JWST Observe First Light and Reionization?



- Detailed hierarchical models (Dr. V. Bromm) show that formation of Pop III stars reionized universe for the first time at  $z \simeq 10-30$  (First Light, age  $\simeq 500-100$  Myr).
- This should be visible to JWST as the first massive stars and surrounding star clusters, and perhaps their extremely luminous supernovae at  $z \simeq 10 \rightarrow 30$ .

We must make sure we theoretically understand the likely Pop III mass-range, their IMF, their duplicity and clustering properties, their SN-rates.


## (5) How will JWST measure First Light & Reionization?



● Can't beat redshift: to see First Light, must observe near-mid IR.

⇒ This is why JWST needs NIRCам at 0.8–5  $\mu\text{m}$  and MIRI at 5–28  $\mu\text{m}$ .

(5) What to expect in Webb (UltraDeep) Fields re. First Light?








841 orbits HUDF 13 filters (false-color): objects affect  $\sim 45\%$  of pixels!!

○  $z=7-8$ , ○  $z=9$ , ○  $z=10-12$ .

592<sup>h</sup> HUDF-weighted log-log: FuvNuvUBViizYJWH, AB  $\lesssim 31$  ( $\gtrsim 2$  nJy).

# HUDF WFC3 IR Galaxy Counts: What to expect in Webb (UltraDeep) Fields?





1.6  $\mu$ m counts (Windhorst<sup>+</sup>2011). [F150W, F225W, F275W, F336W, F435W, F606W, F775W, F850LP, F105W, F125W, F140W not shown].

- Faint-end near-IR count-slope  $\simeq 0.12 \pm 0.02$  dex/mag  $\iff$  Faint-end LF-slope ( $z_{med} \simeq 1.6$ )  $\alpha \simeq -1.4 \Rightarrow$  reach  $M_{AB} \simeq -14$  mag.
- WUDF (- - -) can see  $AB \lesssim 32$  objects:  $M_{AB} \simeq -15$  (LMCs) at  $z \simeq 11$ .
- Lensing will change the landscape for JWST observing strategies.



HST Frontier Field A2744: JWST needs lensing to see First Light at  $z \gtrsim 10-15$ !





Two fundamental limitations determine ultimate JWST image depth:

(1) Cannot-see-the-forest-for-the-trees effect: Background objects blend into foreground neighbors  $\Rightarrow$  Need multi- $\lambda$  deblending algorithms!

(2) House-of-mirrors effect: (Many?) First Light objects can be gravitationally lensed by foreground galaxies  $\Rightarrow$  Must model/correct for this!

● Proper JWST  $2.0\mu\text{m}$  PSF and straylight specs essential to handle this!



## (6) Conclusions

(1) HST set stage to measure galaxy assembly in the last 12.7-13.0 Gyrs.

- Today's Hubble sequence formed 7–10 Gyrs ago.

(2) JWST passed Preliminary & Critical Design Reviews in 2008 & 2010.

Budget and Management replan in 2011. No technical showstoppers!

- More than 80% of JWST H/W built or in fab, & meets/exceeds specs.

(3) JWST is designed to map the epochs of First Light, Reionization, and Galaxy Assembly & SMBH-growth in detail. JWST will determine:

- Formation and evolution of the first star-clusters after 0.2 Gyr.
- How dwarf galaxies formed and reionized the Universe after 1 Gyr.

(4) JWST will have a major impact on astrophysics this decade:

- IR sequel to HST after 2018: Training the next generation researchers.

# SPARE CHARTS

---

- References and other sources of material shown:

<http://www.asu.edu/clas/hst/www/jwst/> [Talk, Movie, Java-tool]

<http://www.asu.edu/clas/hst/www/ahah/> [Hubble at Hyperspeed Java-tool]

<http://www.asu.edu/clas/hst/www/jwst/clickonHUDF/> [Clickable HUDF map]

<http://www.jwst.nasa.gov/> & <http://www.stsci.edu/jwst/>

<http://ircamera.as.arizona.edu/nircam/>

<http://ircamera.as.arizona.edu/MIRI/>

<http://www.stsci.edu/jwst/instruments/nirspec/>

<http://www.stsci.edu/jwst/instruments/fgs>

Gardner, J. P., et al. 2006, Space Science Reviews, 123, 485–606

Mather, J., & Stockman, H. 2000, Proc. SPIE Vol. 4013, 2

Windhorst, R., et al. 2008, Advances in Space Research, 41, 1965


Windhorst, R., et al., 2011, ApJS, 193, 27 (astro-ph/1005.2776)

## (4b) Predicted Galaxy Appearance for JWST at redshifts $z \simeq 1-15$

NGC 3310

ESO0418-008

UGC06471-2



**Ultraviolet Galaxies**


**HST • WFPC2**

NASA and R. Windhorst (Arizona State University) • STScI-PRC01-04

- The rest-frame UV-morphology of galaxies is dominated by young and hot stars, with often significant dust imprinted (Mager-Taylor et al. 2005).
- High-resolution HST ultraviolet images are benchmarks for comparison with very high redshift galaxies seen by JWST.

## (4b) Predicted Galaxy Appearance for JWST at redshifts $z \simeq 1-15$

HST  $z=0$  JWST  $z=2$   $z=5$   $z=9$   $z=15$



With Hubble UV-optical images as benchmarks, JWST can measure the evolution of galaxy structure & physical properties over a wide range of cosmic time:

- (1) Most spiral disks will dim away at high redshift, but most formed at  $z \lesssim 1-2$ .

Visible to JWST at very high  $z$  are:

- (2) Compact star-forming objects (dwarf galaxies).
- (3) Point sources (QSOs).
- (4) Compact mergers & train-wrecks.

# Northrop Grumman Expertise in Space Deployable Systems

- Over 45 years experience in the design, manufacture, integration, verification and flight operation of spacecraft deployables
- 100% mission success rate, comprising over 640 deployable systems with over 2000 elements





### Baseline "Cup Down" Tower Configuration at JSC (Before)



### JSC "Cup Up" Test Configuration (New Proposal)




Most recent Tower Design shows an Inner Optical Tower supported by a Outer structure with Vibration Isolation at the midplane. Everything shown is in the 20K region (helium connections, etc. not shown) except clean room and lift fixture.

Current plan calls for 33KW cooldown capability, 12 KW steady state, 300-500mW N2 cooling

JSC currently has 7 KW He capability


Current plan includes 10 trucks of LN2/day during cooldown

Interferometers, Sources, Null Lens and Alignment Equipment Are in Upper and Lower Pressure Tight Enclosure Inside of Shroud



No Metrology Tower and Associated Cooling H/W. External Metrology  
Two basic test options:  
1. Use isolators, remove drift through fast active control + freeze test equipment jitter  
2. Eliminate vibration isolators (but use soft dampeners) to avoid drift, freeze out jitter  
Builds on successful AMSD heritage of freezing and averaging jitter, testing through windows.


Possible payload "floor" to separate ambient pressure and temperature.



Drawing care of ITT


- JWST underwent several significant replans and risk-reduction schemes:**
- $\lesssim 2003$ : Reduction from 8.0 to 7.0 to 6.5 meter. Ariane-V launch vehicle.
  - 2005: Eliminate costly 0.7-1.0  $\mu\text{m}$  performance specs (kept 2.0  $\mu\text{m}$ ).
  - 2005: Simplification of thermal vacuum tests: cup-up, not cup-down.
  - 2006: All critical technology at Technical Readiness Level 6 (TRL-6).
  - 2007: Further simplification of sun-shield and end-to-end testing.
  - 2008: Passes Mission Preliminary Design & Non-advocate Reviews.
  - 2011: Passes Mission Critical Design Review — Replan Int. & Testing.

**First light  
NIRCam**



1.  
Segment  
Image  
Capture

**After Step 1**



**Initial Capture**

18 individual 1.6-m diameter aberrated sub-telescope images  
 PM segments: < 1 mm, < 2 arcmin tilt  
 SM: < 3 mm, < 5 arcmin tilt


**Final Condition**

PM segments:  
 < 100  $\mu\text{m}$ ,  
 < 2 arcsec tilt  
 SM: < 3 mm,  
 < 5 arcmin tilt

**2. Coarse Alignment**

Secondary mirror aligned  
 Primary RoC adjusted

**After Step 2**




Primary Mirror segments:  
 < 1 mm, < 10 arcsec tilt  
 Secondary Mirror :  
 < 3 mm, < 5 arcmin tilt

WFE < 200  $\mu\text{m}$  (rms)

**3. Coarse Phasing - Fine Guiding (PMSA piston)**

**After Step 3**




WFE: < 250  $\mu\text{m}$  rms

WFE < 1  $\mu\text{m}$  (rms)

**4. Fine Phasing**

**After Step 4**



WFE: < 5  $\mu\text{m}$  (rms)

WFE < 110 nm (rms)

**5. Image-Based Wavefront Monitoring**

**After Step 5**




WFE: < 150 nm (rms)

WFE < 110 nm (rms)


JWST's Wave Front Sensing and Control is similar to the Keck telescope.

In L2, need WFS updates every 10 days depending on scheduling/illumination.






Wave-Front Sensing tested hands-off at 40 K in 1-G at JSC in 2015-2016.  
Ball 1/6 scale-model for WFS: produces diffraction-limited  $2.0 \mu\text{m}$  images.



JWST can observe NEP+SEP continuously: Think of 1000-hr proposals!

V3 (anti-spacecraft)



OTE ISIM



(V1, V3)  
origin

Tertiary  
Mirror

Secondary mirror

Cassegrain  
focus

Fine  
Steering Mirror


Focal  
Surface

Primary  
Mirror

f/#: 20.0

Effective Focal Length: 131.4 m

PM diameter = 6.6 m (circumscribed circle)



# One day we will need a UV-optical sequel to Hubble:





Figure 2.5.3 - HORUS Observatory and Optical Ray Trace



[Left] One of two spare 2.4 m NRO mirrors: one will become WFIRST.

- NASA may look for partners to turn 2nd NRO into UV-opt HST sequel.

[Middle] HORUS: 3-mirror anastigmat NRO as UV-opt HST sequel.

- Can do wide-field ( $\sim 0.25$  deg) UV-opt  $0''.06$  FWHM imaging to  $AB \lesssim 30$  mag, and high sensitivity (on-axis) UV-spectroscopy (Scowen et al. 2012).

[Right] ATLAST: 8–16 m UV-opt HST sequel, with JWST heritage.

- Can do same at 9 m.a.s. FWHM routinely to  $AB \lesssim 32-34$  mag, [and an ATLAST-UDF to  $AB \lesssim 38$  mag  $\sim 1$  pico-Jy].



Life-sized JWST model, at NASA/GSFC with the whole JWST Project ...



Life-sized JWST model, at NASA/GSFC Friday afternoon after 5 pm ...