


Reionization at $z \simeq 2-6$ — Whose dunnit: Dwarfs or Giants?

Rogier Windhorst (ASU) — JWST Interdisciplinary Scientist

Brent Smith, Junehyoung Jeon (UG!) S. Cohen, R. Jansen, L. Jiang, M. Dijkstra,

A. Inoue, A. Koekemoer, R. Bielby, J. MacKenty, R. O'Connell, & J. Silk






Summary Talk Princeton Bahcall Lunch, Princeton (NJ; via Zoom); Tu. Nov. 10, 2020

Talk is on: http://www.asu.edu/clas/hst/www/jwst/jwsttalks/princeton20_Bahcall_lunch.pdf





Answer: Weak AGN at $z \simeq 2.3-3.5$, but galaxies with $\langle f_{esc} \rangle \simeq 0.4$ at $z \simeq 6$

HST WFC3 UVIS image stacking of galaxies & weak AGN at $z \simeq 2.3-3.5$:



B. Smith et al. (2018, ApJ, 853, 191; 2020 ApJ, 897, 41):




- Weak AGN (1 out of 17) dominate reionizing flux at $z \simeq 2.3-3.5$!
- Faint galaxies (94 objects) in GOODS-N+S contribute marginal LyC flux.



- One lucky AGN has $f_{esc} \simeq 24 \pm 4\%$, higher than most galaxies.
- Ensemble LyC of 111 objects has $f_{esc} \simeq 3-30\%$ at $z \simeq 2.3-3$.
- Large (IGM τ induced) errors, but f_{esc} trending higher with redshift?




ID63 $z=6.027$ $\chi^2=0.99$ $Z=0.004$ Age=350.0 $f_{esc}=0.0$ $\beta=-2.43$

ID62 $z=6.964$ $\chi^2=1.21$ $Z=0.004$ Age=400.0 $f_{esc}=1.0$ $\beta=-2.5$



J. Jeon et al. (2020, astro-ph/2011; AJ subm): 47 SDF Ly α galaxies at $z \simeq 6$:

● CIGALE fits of Subaru-HSC/WFC3-IR/Spitzer SEDs: $\langle f_{esc} \rangle \simeq 0.4$ at $z \simeq 6$.



● CIGALE model E_{B-V} vs. UV-slope offset from Calzetti/Meurer $z \simeq 0$ relation: JWST NIRSpec targets!

● f_{esc} may trend *higher* for redder β -slopes and higher E_{B-V} (!): Older, dustier SEDs have more ISM holes?